

### Behavioral Health HEDIS® Measure Overview

Measure	Patient Population	Multiple Numerators	Compliance Requirements	Details*
Follow-Up Care for Children Prescribed ADHD Medication <u>ADD-E</u>	Newly prescribed ADHD medication.	<b>Initiation Phase</b> 30 days of when ADHD medication was first dispensed. <b>Continuation and Maintenance Phase</b> 270 days following Initiation Phase	<ul style="list-style-type: none"> <li>1 visit within 30 days of when medication was first dispensed (initiation).</li> <li>At least 2 follow-up visits within 270 days with <b>ANY practitioner</b> (9 months) after the initiation phase ended. (Continuation).</li> </ul>	Age: 6-12 Telehealth acceptable: Yes
Antidepressant Medication Management <u>AMM</u>	Major depression diagnosis and antidepressant medication management.	<b>Effective Acute Phase</b> Treatment (84d, 12w) <b>Effective Continuation Phase</b> Treatment (180d, 6m)	<ul style="list-style-type: none"> <li>Remained on an antidepressant medication for at least 84 days or 12 weeks (Acute).</li> <li>Remained on antidepressant medication for at least 180 days or 6 months (Continuation).</li> </ul>	Age: 18+ Telehealth acceptable: Yes
Metabolic Monitoring for Children and Adolescents on Antipsychotics <u>APM-E</u>	Dispensed 2 or more anti-psychotic prescriptions.		<ul style="list-style-type: none"> <li>Metabolic testing (blood glucose and cholesterol) within measurement year.</li> </ul>	Age: 1-17
Use of First-Line Psychosocial Care for Children and Adolescents on Antipsychotics <u>APP</u>	Newly prescribed anti-psychotic medication dispensed <u>after</u> psychosocial care.		<ul style="list-style-type: none"> <li>Documentation of <b>psychosocial care as first-line treatment</b>.</li> <li>Negative medication history (120 days prior to dispensed medication).</li> </ul>	Age: 1-17 Telehealth acceptable: Yes
Depression Screening and Follow-Up for Adolescents and Adults <u>DSF-E</u>	Members 12 and older that have a positive screening for depression.		<ul style="list-style-type: none"> <li>Clinical depression screening using a standardized tool.</li> <li>Follow-up on a <b>positive</b> screen within 30 days with <b>ANY practitioner</b>.</li> </ul>	Age: 12+ Telehealth acceptable: Yes
Follow-Up After Emergency Department Visit for Substance Use <u>FUA</u>	ED visit with a primary diagnosis of substance use disorder (SUD), or any diagnosis of drug overdose.	7/ 30-day follow-up	<ul style="list-style-type: none"> <li>Follow-up with <b>ANY practitioner</b> within 7 days/30 days of ED visit with a diagnosis of substance use disorder (SUD) or drug overdose.</li> </ul>	Age: 13+ Telehealth acceptable: Yes
Follow-Up After Hospitalization for Mental Illness <u>FUH</u>	Hospitalization for a principal diagnosis of mental illness/intentional self-harm.	7/ 30-day follow-up	<ul style="list-style-type: none"> <li>Follow-up with a <b>mental health provider</b> within 7 days/30 days of discharge.</li> </ul>	Age: 6+ Telehealth acceptable: Yes Mental health provider required: Yes
Follow-Up After High-Intensity Care for Substance Use Disorder <u>FUI</u>	Members with a principal diagnosis of substance use disorder (SUD) from inpatient hospitalization, residential treatment or withdrawal management.	7/ 30-day follow-up	<ul style="list-style-type: none"> <li>Follow-up with <b>ANY practitioner</b> within 7 days/30 days of ED visit with a principal diagnosis of substance use disorder (SUD) or drug overdose.</li> </ul>	Age: 13+ Telehealth acceptable: Yes Principal diagnosis required: Yes

\*Reported for both commercial & Medicaid populations unless otherwise indicated.

### Behavioral Health HEDIS® Measure Overview

Measure	Patient Population	Multiple Numerators	Compliance Requirements	Details*
Follow-Up After Emergency Department Visit for Mental Illness <u>FUM</u>	ED visit with principal diagnosis of mental illness/intentional self-harm.	7/ 30-day follow-up	<ul style="list-style-type: none"> <li>Follow-up with <b>ANY practitioner</b> within 7 days/ 30 days of ED visit with a principal diagnosis of a mental health disorder <b>or</b> intentional self-harm and any diagnosis of a mental health disorder.</li> </ul>	Age: 6+ Telehealth acceptable: Yes Principal diagnosis required: Yes
Initiation and Engagement of Substance Use Disorder Treatment <u>IET</u>	New substance use disorder (SUD) episodes in any setting.	<b>Initiation:</b> New SUD episodes with treatment within 14 days <b>Engagement:</b> New SUD episodes with evidence of treatment engagement within 34 days of initiation.	<ul style="list-style-type: none"> <li>Initiation of SUD treatment within 14 days of episode (inpatient, outpatient, intensive outpatient, partial hospitalization, telehealth, medication treatment). One treatment visit.</li> <li>Engagement of SUD treatment within 34 days of initiation. Two treatment visits.</li> </ul>	Age: 13+ Telehealth acceptable: Yes
Pharmacotherapy for Opioid Use Disorder <u>POD</u>	New opioid use disorder (OUD) pharmacotherapy events.		<ul style="list-style-type: none"> <li>OUD pharmacotherapy events lasting at least 180 days.</li> </ul>	Age: 16+
Adherence to Antipsychotic Medications for Individuals with Schizophrenia <u>SAA</u>	Schizophrenia or schizoaffective disorder dispensed anti-psychotic medication.		<ul style="list-style-type: none"> <li>Remained on anti-psychotic medication for at least 80% of the treatment period.</li> </ul>	Age: 18+ Telehealth acceptable: Yes
Cardiovascular Monitoring for People with Cardiovascular Disease and Schizophrenia <u>SMC</u>	Schizophrenia or schizoaffective disorder and cardiovascular disease.		<ul style="list-style-type: none"> <li>LDL-C testing during the measurement year.</li> </ul>	Age: 18-64 Reporting Population: Medicaid only
Diabetes Monitoring for People with Diabetes and Schizophrenia <u>SMD</u>	Schizophrenia or schizoaffective disorder and diabetes		<ul style="list-style-type: none"> <li>HbA1c <b>AND</b> LDL-C testing during the measurement year.</li> </ul>	Age: 18-64 Reporting Population: Medicaid only
Diabetes Screening for People with Schizophrenia or Bipolar Disorder Who Are Using Antipsychotic Medications <u>SSD</u>	Schizophrenia, schizoaffective disorder or bipolar disorder with antipsychotic medication dispensed during measurement year.		<ul style="list-style-type: none"> <li>Diabetes testing during the measurement year (HbA1c <b>OR</b> glucose test).</li> </ul>	Age: 18-64 Reporting Population: Medicaid only

\*Reported for both commercial & Medicaid populations unless otherwise indicated.  
 This is not an exhaustive list of criteria as coding standardization and other behavioral health practices can impact compliance.

For more information scan the QR code or visit: <https://healthplanofnevada.com/provider/hedis-measures>

